

A Review of CAS Mathematical Capabilities

Michael Wester
Cotopaxi, 1801 Quincy, SE
Albuquerque, New Mexico 87108, USA
`wester@math.unm.edu`

November 17, 1994

Abstract

Computer algebra systems (CASs) have become an important computational tool in the last decade. General purpose CASs, which are designed to solve a wide variety of problems, have gained special prominence. In this paper, the capabilities of six major general purpose CASs (Axiom, Derive, Macsyma, Maple, Mathematica and Reduce) are reviewed on 131 short problems covering a broad range of (primarily) symbolic mathematics.

A demo was developed for each CAS, run and the results evaluated. Problems were graded in terms of whether it was easy or difficult or possible to produce an answer and if an answer was produced, whether it was correct. It is the author's hope that this review will encourage the development of a comprehensive CAS test suite.

Presented below is a summary of 131 mathematical problems (primarily symbolic) that were given to the six general purpose computer algebra systems (CASs) listed in Table 1. The CAS versions tested were those that were available to the author and were typically the newest and most comprehensive versions that were generally available at the time of this evaluation.

The notations used in the summary are explained in Table 2. A demo was developed for each CAS, run and the results evaluated. Problems easily and successfully solved are marked by a ●. Those that required more effort than a mere 'simplify' (sometimes considerably more effort) or the results could not be completely simplified or were incomplete in some way are marked with an ○. If the problem could not be solved by the CAS, the corresponding entry in the summary table was left blank. Incorrect answers are indicated by an ×.

Problem descriptions are abbreviated due to space limitations with this format and may not always be complete. However, the complete demos and their output are readily obtained by anonymous FTP from `math.unm.edu`. The directory `pub/cas` will contain current versions of these and other potentially interesting CAS related files.

The philosophy that I followed in making these comparisons included several facets. The choice of problems was deliberately broad as a primary goal was to provide a useful indication of the breadth of coverage of each general purpose system. A secondary goal was to give a

feeling of the depth of coverage provided for certain classes of problems. This aspect of the survey is certainly incomplete as right now some areas of mathematics are covered by a more interesting range of examples than others (e.g., the problems for matrix algebra, products and limits currently are all quite easy). It is my intention to remedy this unevenness in a future version of this review.

The primary emphasis of this review is on exact, symbolic mathematics, although a few approximate, numerical problems are also included. I have totally ignored for the time being issues of graphics, language design, user interface and computational speed, all of which are important as well. The six general purpose CASs examined can, with varying amounts of user assistance, solve a great variety of problems. I have tried to emphasize those problems that ideally should involve minimal user intervention.

If it was necessary to tell a system most of the steps to solve a problem, I did not consider this a solution by the CAS.¹ Poorly documented methods and results were also judged unfavorably. If only ‘some’ user intervention was required, this was acknowledged by an ○. If at most minimal aid was provided by the user, only then was a ● rewarded.² Packages not provided with the standard distribution of a CAS were excluded from consideration. Many systems do, however, have some quite nice user packages that are only available (easily) on the Internet/via email or must be purchased separately.

Some final comments: in order to perform this review, it was necessary to dig deeply into each of the CASs’ reference manuals. None of them were particularly easy to use without a great deal of study and unfortunately, the indices in general were of fairly limited utility (Mathematica’s was the best of the lot). Other people have also reviewed the software in or made comparisons of multiple CASs, emphasizing different aspects than was done here. For example, see [Har91, Her94, Sim92]. In summary, the general purpose CASs surveyed here are all quite powerful, each with its own particular strengths and weaknesses. This review should only be used as a guide, not as a definitive basis for comparison. Comments are welcome and additional problems are actively solicited.

References

- [Asl94] Helmer Aslaksen, *Multiple-valued complex functions and computer algebra*, Research Report No. 631, Department of Mathematics, National University of Singapore, October 1994.
- [Har91] David Harper, Chris Wooff and David Hodgkinson, *A Guide to COMPUTER ALGEBRA SYSTEMS*, John Wiley & Sons, 1991.
- [Her94] W. Hereman, “Review of Symbolic Software for the Computation of Lie Symmetries of Differential Equations”, *Euromath Bulletin*, Volume 1, Number 2, 1994, 45–79.
- [Rob93] Nicolas Robidoux, “Does Axiom Solve Systems of O.D.E.’s Like Mathematica?”, LA-UR-93-2235, Los Alamos National Laboratory, Los Alamos, New Mexico.

¹However, after receiving some artful responses from various CAS developers, I may institute a ‘super tricky’ category in a future version of this review!

²Note: the complex domain problems, #37–45, were strictly evaluated for consistency with the domains assumed by the variables involved—see also [Asl94].

[Sim92] Barry Simon, “Comparative CAS Reviews”, *Notices of the American Mathematical Society*, Volume 39, Number 7, September 1992, 700–710.

Acknowledgements

Paul Abbott	(Mathematica)
Richard J. Fateman	(Macsyma)
Jeffrey P. Golden	(Macsyma)
Tony Hearn	(for making Reduce 3.5 available)
Tom Issaevitch	(Mathematica)
Richard Liska	(Reduce)
Rolf Mertig	(Mathematica)
Michael Monagan	(Maple)
Albert D. Rich	(for making Derive XM Version 3 available)
Nicolas Robidoux	(Axiom, triangular system of two ODEs example [Rob93])
Margaret Sanders	(statistics examples)
Stanly Steinberg	(several examples)
David Wood	(readability)

Footnotes to the Table of Problems

- ¹ biased rather than unbiased estimator used (division by n rather than by $n - 1$)
- ² pattern matching is primitive
- ³ `evalc` is correct, but `simplify` is incorrect
- ⁴ this happens automatically!
- ⁵ produces complete general solution!
- ⁶ one of the answers produced is incorrect
- ⁷ numerical evaluation of ζ worked only with big floats in this version
- ⁸ lack of documentation on the answer that is returned
- ⁹ incorrect answer produced with `noPole` option
- ¹⁰ ignores assumptions on a
- ¹¹ limiting the order to 0 or 1 (Maple) or 1 or 2 (Mathematica) produces errors
- ¹² answer contains an unevaluated (but obvious) limit
- ¹³ requires some user sophistication to solve
- ¹⁴ claims that this is a partial differential equation!
- ¹⁵ produces only the trivial solution
- ¹⁶ need to be careful about the statement order
- ¹⁷ definition is ‘hidden’ within one of the standard packages

Axi	Axiom 1.2	March 17, 1993
Der	Derive XM Version 3	1994
Mac	Macsyma 419.0	1994
Mpl	Maple V Release 3	1994
Mma	Mathematica 2.2	1993
Red	Reduce 3.5	15-Oct-93

Table 1 CASs used.

•	success! (hurrah)
◦	success, but indirectly, incomplete or unsimplified could not do the problem (boo)
×	produced the wrong answer (hiss)
⊗	partial success, but also partially incorrect (hmmm)
⇒	yields
→	then
"	previous result
D	derivative operator
diff	differentiate
factor(..., α)	factor over the algebraic field extension α
Im z	imaginary part of z
$J_\mu(z)$	Bessel function of the first kind of order μ
$K^h_{i\ jk \ell}$	covariant derivative of the Riemann curvature tensor
laplace($f(t)$, $t \rightarrow s$)	Laplace transform of $f(t)$ under $t \rightarrow s$
N(..., $\{, k\}$)	numerically evaluate (to k -digit precision)
(operator)	define as an operator
pade($f(x)$, $x = a$)	Pade approximation of $f(x)$ about $x = a$
poly	polynomial
power_series($f(x)$, $x = a$)	general power series formula of $f(x)$ about $x = a$
Re z	real part of z
(real)	produce an explicitly real solution
rectform($f(z)$)	rectangular form of the complex function $f(z)$
(rewrite rules)	using user supplied rewrite rules
solve($x = f(y)$, $y = a$, series)	solve for $y(x)$ about $y = a$ using series reversion
stdev(...)	unbiased sample standard deviation
taylor($f(x)$, $x = a$)	truncated Taylor series of $f(x)$ about $x = a$
$\zeta(n)$	Riemann zeta function

Table 2 Notations used.

#	PROBLEM	Axi	Der	Mac	Mpl	Mma	Red
1	50!	•	•	•	•	•	•
2	<code>factor(50!) $\Rightarrow 2^{47}3^{22}5^{12}7^811^413^3 \dots 47$</code>	•	•	•	•	•	○
3	$\frac{1}{2} + \dots + \frac{1}{10} \Rightarrow \frac{4861}{2520}$	•	•	•	•	•	•
4	$N(e^{\pi\sqrt{163}}, 50) \approx 262537412640768744.0$	•	•	•	•	•	•
5	$N(J_2(1+i)) \approx 0.04158 + 0.24740i$	•	•	•		•	•
6	$N(\frac{1}{7}) \Rightarrow 0.142857$	•					
7	continued fraction of 3.1415926535	•		•	•	•	
8	$\sqrt{2\sqrt{3}+4} \Rightarrow 1 + \sqrt{3}$		•	•	•		•
9	$\sqrt{14+3\sqrt{3}+\dots} \Rightarrow 3 + \sqrt{2}$		•	○	•		
10	$2\infty - 3 \Rightarrow \infty$	•	•	○	○	•	
11	<code>stdev([1,2,3,4,5]) $\Rightarrow \sqrt{\frac{5}{2}}$</code>		○ ¹	•	•	•	
12	hypothesis testing: t distribution		× ¹	○	○	•	
13	hypothesis testing: normal distribution		○ ¹	○	○	•	
14	$\frac{x^2-4}{x^2+4x+4} \Rightarrow \frac{x-2}{x+2}$	•	•	•	•	•	•
15	$\frac{e^x-1}{e^{x/2}+1} \Rightarrow e^{x/2} - 1$	•	•	•	○	•	
16	<code>expand((x+1)²⁰) \rightarrow diff \rightarrow factor</code>	•	•	•	•	•	•
17	<code>factor($x^{100} - 1$)</code>	•	•	•	•	•	•
18	<code>factor($x^4 - 3x^2 + 1$, RootOf($\phi^2 - \phi - 1$))</code>	•		•	•		
19	<code>factor($x^4 - 3x^2 + 1$) mod 5</code>	•		•	•	•	•
20	$\frac{x^2+2x+3}{x^3+4x^2+5x+2} \Rightarrow \frac{3}{x+2} - \frac{2}{x+1} + \frac{2}{(x+1)^2}$	○	•	•	•	•	•
21	<code>assume($x \geq y, y \geq z, z \geq x$); is($x = z$)</code>			•	×		
22	<code>assume($x > y, y > 0$); is($2x^2 > 2y^2$)</code>			•	•		
23	<code>solve($x-1 > 2$) $\Rightarrow x < -1$ or $x > 3$</code>		•		•		
24	<code>solve(expand(($x-1$)\dots($x-5$)) < 0)</code>		•		•		
25	$\frac{\cos 3x}{\cos x} \Rightarrow \cos^2 x - 3\sin^2 x$ (or similar)	○	•	•	•	•	
26	$\frac{\cos 3x}{\cos x} \Rightarrow 2\cos 2x - 1$	○	•	•	•	•	
27	$\frac{\cos 3x}{\cos x} \Rightarrow \cos^2 x - 3\sin^2 x$ (rewrite rules)	•		•	○ ²	•	•
28	$\sqrt{997} - (997^3)^{1/6} \Rightarrow 0$	•	•	•	•	•	○
29	$\sqrt{999983} - (999983^3)^{1/6} \Rightarrow 0$	•	•	•	•		○
30	$(2^{1/3} + 4^{1/3})^3 - 6(2^{1/3} + 4^{1/3}) - 6 \Rightarrow 0$		•	•	•	•	•
31	$\log(\tan(\frac{1}{2}x + \frac{\pi}{4})) - \sinh^{-1}(\tan x) \Rightarrow 0$			○	○		
32	derivative of above is 0 & above at 0 is 0	○	○	•	•	○	
33	$\log \frac{2\sqrt{r+1}}{\sqrt{4r+4}\sqrt{r+1}} \Rightarrow 0$		•	•		•	•

#	PROBLEM	Axi	Der	Mac	Mpl	Mma	Red
34	$(4r + 4\sqrt{r} + 1)^{\frac{r}{2\sqrt{r}+1}}(2\sqrt{r} + 1)^{\frac{1}{2\sqrt{r}+1}} \dots$	●	●			○	
35	$\text{rectform}(\log(3 + 4i)) \Rightarrow \log 5 + i \tan^{-1} \frac{4}{3}$	○	○	●	●	○	
36	$\text{rectform}(\tan(x + iy))$	○	○	●	●	●	
37	$\frac{\sqrt{xy z ^2}}{\sqrt{x z }} \Rightarrow \frac{\sqrt{xy}}{\sqrt{x}} \not\Rightarrow \sqrt{y}$	○	●	●	●	○	×
38	$\sqrt{\frac{1}{z} - \frac{1}{\bar{z}}} \Rightarrow 0 \quad (z \text{ is not real negative})$	×	●	●	●	●	×
39	$\sqrt{e^z} - e^{z/2} \Rightarrow 0 \quad (-\pi < \text{Im } z \leq \pi)$	○	●	●	×	○	×
40	$\sqrt{e^{6i}} \Rightarrow -e^{3i} \quad (\text{principal value})$		○	○	⊗ ³	○	×
41	$\log e^z \Rightarrow z \quad (-\pi < \text{Im } z \leq \pi)$	●	●	×	○	●	×
42	$\log e^{10i} \Rightarrow (10 - 4\pi)i \quad (\text{principal value})$		●		○	○	
43	$(xy)^{1/n} - x^{1/n}y^{1/n} \Rightarrow 0 \quad (\text{Re } x, \text{Re } y > 0)$	×	●	●	●	●	×
44	$\tan^{-1}(\tan z) \Rightarrow z \quad (-\frac{\pi}{2} < z \leq \frac{\pi}{2})$	×	○	○	○	●	●
45	$\tan^{-1}(\tan 4) \Rightarrow 4 - \pi \quad (\text{principal value})$		●	○	●	○	
46	$\frac{x=0}{2} + 1 \Rightarrow \frac{x}{2} + 1 = 1$		●	●	●	×	×
47	$\text{solve}(3x^3 - 18x^2 + 33x - 19 = 0) \quad (\text{real})$		● ⁴	●	●	●	● ⁴
48	$\text{solve}(x^4 + x^3 + x^2 + x + 1 = 0)$	○	●	●	●	●	●
49	verify a solution of the above	●	●	●	●	●	●
50	$\text{solve}(e^{2x} + 2e^x + 1 = z, x)$	●	●	●	●	●	● ⁵
51	$\text{solve}((x + 1)(\sin^2 x + 1)^2 \cos^3 3x = 0)$		○	○	○	○	● ⁵
52	$\text{solve}(e^z = 1) \Rightarrow z = 0 [+n2\pi i]$	●	●	●	●	●	● ⁵
53	$\text{solve}(\sin x = \cos x) \Rightarrow x = \frac{\pi}{4} [+n\pi]$	●	●		●		○
54	$\text{solve}(\tan x = 1) \Rightarrow x = \frac{\pi}{4} [+n\pi]$	●	●	●	●	●	● ⁵
55	$\text{solve}(\sin x = \tan x) \Rightarrow 0, 0 [+n\pi, +n2\pi]$	○	○	●	●		○
56	$\text{solve}(\sqrt{x^2 + 1} = x - 2) \Rightarrow x = \{\}$	×	○		○	●	×
57	$\text{solve}(e^{2-x^2} = e^{-x}) \Rightarrow x = \{-1, 2\}$		●	●	○		
58	$\text{solve}(\sqrt{\log x} = \log \sqrt{x}) \Rightarrow x = \{1, e^4\}$	⊗ ⁶			●		
59	$\text{solve}(x - 1 = 2) \Rightarrow x = \{-1, 3\}$		●	●	●		●
60	solve a 3×3 dependent linear system	●	●	●	●	●	●
61	solve a system of nonlinear equations	●		●	●	●	●
62	invert a 2×2 symbolic matrix	●	●	●	●	●	●
63	$\det(4 \times 4 \text{ Vandermonde matrix})$	●	●	●	●	●	●
64	eigenvalues of a 3×3 integer matrix	●	●	●	●	●	●
65	tensor covariant derivative			●			
66	$K_i^h{}_{jk l} + K_i^h{}_{kl j} + K_i^h{}_{lj k} \Rightarrow 0 \quad (\text{Bianchi})$			●			

#	PROBLEM	Axi	Der	Mac	Mpl	Mma	Red
67	$\sum_{k=1}^n k^3 \Rightarrow \frac{n^2(n+1)^2}{4}$	•	•	•	•	•	•
68	$\sum_{k=1}^{\infty} (\frac{1}{k^2} + \frac{1}{k^3}) \Rightarrow \frac{\pi^2}{6} + \zeta(3)$		•	•	•	•	
69	$N(\sum_{k=1}^{\infty} (\frac{1}{k^2} + \frac{1}{k^3})) \approx 2.84699$		•	• ⁷	•	•	
70	$\prod_{k=1}^n k \Rightarrow n!$		•	•	•	•	
71	$\lim_{n \rightarrow \infty} (1 + \frac{1}{n})^n \Rightarrow e; \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} \Rightarrow \frac{1}{2}$	•	•	•	•	•	•
72	$\frac{d^2}{dt^2} y(x(t)) \Rightarrow \frac{d^2 y}{dx^2} \left(\frac{dx}{dt}\right)^2 + \frac{dy}{dx} \frac{d^2 x}{dt^2}$	•	○	•	•	•	
73	$\int \frac{1}{x^3+2} dx \rightarrow \text{diff} \rightarrow \text{simplify}$	•	•	•	•	•	•
74	$\int \frac{1}{a+b \cos x} dx \quad (a < b)$	•		•	•		
75	$\frac{d}{dx} \int \frac{1}{a+b \cos x} dx = \frac{1}{a+b \cos x}$	•	•	•	•	•	○
76	$\frac{d}{dx} x \Rightarrow \frac{x}{ x } \text{ or } \text{sign}(x)$	•	•	•	○ ⁸	○	•
77	$\int x dx \Rightarrow \frac{x x }{2} \quad (\text{for real } x)$		•	•	○	•	•
78	$\frac{d}{dx} x \quad (\text{piecewise defined})$	×	×	•	•	•	
79	$\int x dx \quad (\text{piecewise defined})$	×		•			
80	$\int \frac{x}{\sqrt{1+x} + \sqrt{1-x}} dx \Rightarrow \frac{(1+x)^{3/2} + (1-x)^{3/2}}{3}$	•	•	○	•	•	•
81	$\int \frac{\sqrt{1+x} - \sqrt{1-x}}{2} dx \Rightarrow \frac{(1+x)^{3/2} + (1-x)^{3/2}}{3}$	•	•	•	•	•	•
82	$\int_{a-1}^{a+1} \frac{1}{x-a} dx \Rightarrow 0 \quad (\text{principal value})$		×	•	×	×	
83	$\int_{a-1}^{a+1} \frac{1}{(x-a)^2} dx \Rightarrow \text{divergent}$		×	•	×	×	
84	$\int_0^1 \sqrt{x + \frac{1}{x} - 2} dx \Rightarrow \frac{4}{3}$	⁹	•	•	•	•	
85	$\int_1^2 \sqrt{x + \frac{1}{x} - 2} dx \Rightarrow \frac{4-\sqrt{8}}{3}$	○	•	•	×	•	
86	$\int_0^2 \sqrt{x + \frac{1}{x} - 2} dx \Rightarrow \frac{8-\sqrt{8}}{3}$	⁹	•		×	•	
87	$\int_{-\infty}^{\infty} \frac{\cos x}{x^2+a^2} dx \quad (a > 0) \Rightarrow \frac{\pi}{a} e^{-a}$			•	×	○ ¹⁰	
88	$\int_0^{\infty} \frac{t^{a-1}}{1+t} dt \quad (0 < a < 1) \Rightarrow \frac{\pi}{\sin \pi a}$			•	○ ¹⁰	• ¹⁰	
89	$\int_0^{\infty} \left(\frac{J_1(x)}{x}\right)^2 dx \Rightarrow \frac{4}{3\pi}$				×	•	
90	$\int_0^a \int_0^{b(1-x/a)} \int_0^{c(1-x/a-y/b)} 1 dz dy dx \Rightarrow \frac{abc}{6}$	•	•	•	•	•	
91	$\text{taylor}(\frac{1}{\sqrt{1-(v/c)^2}}, v = 0)$	•	•	•	•	•	•
92	$\frac{1}{\text{above}^2} \Rightarrow 1 - \frac{v^2}{c^2} + \dots$	•	○	•	○	•	○
93	$\frac{\text{taylor}(\sin x, x=0)}{\text{taylor}(\cos x, x=0)} = \text{taylor}(\tan x, x = 0)$	•	○	•	○	•	○
94	$\text{taylor}((\log x)^a e^{-bx}, x = 1)$			•			○
95	$\text{taylor}(\log(\sinh z) + \log(\cosh(z + w)))$			•	○ ¹¹	○ ¹¹	•
96	" - $\text{taylor}(\log(\sinh z \cosh(z + w)), z = 0)$			•	○ ¹¹	○ ¹¹	•
97	$\text{taylor}(\log(\frac{\sin x}{x}), x = 0)$	•	•	•	•	•	•
98	$\text{power_series}(\log(\frac{\sin x}{x}), x = 0)$			•			
99	$\text{power_series}(e^{-x} \sin x, x = 0)$			○	•		

#	PROBLEM	Axi	Der	Mac	Mpl	Mma	Red
100	$\text{solve}(x = \sin y + \cos y, y = 0, \text{series})$			•	•	•	
101	$\text{pade}(e^{-x}, x = 0) \Rightarrow \frac{2-x}{2+x}$	•	•	○	•	•	
102	$\text{laplace}(\cos((\omega - 1)t), t \rightarrow s) \Rightarrow \frac{s}{s^2 + (\omega - 1)^2}$	•	○ ¹²	•	•	•	•
103	inverse Laplace transform of above	•		•	•	•	○
104	$\text{solve}([r_{n+2} - 2r_{n+1} + r_n = 2, \dots], r_n)$		○	○	•	•	
105	$\text{solve}([\frac{d^2 f}{dt^2} + 4f = \sin 2t, f(0) = f'(0) = 0])$	•	○	•	○	•	○
106	above solution using Laplace transforms			•	•	•	
107	$\text{solve}(x^2 \frac{dy}{dx} + 3xy = \frac{\sin x}{x}, y(x))$	•	○	•	•	•	•
108	$\text{solve}(\frac{d^2 y}{dx^2} + y(\frac{dy}{dx})^3 = 0, y(x))$		○ ¹³	•	•		
109	$\text{solve}(\frac{d}{dx}y(x, a) = ay(x, a), y(x, a))$			•	14	14	•
110	$\text{solve}([\frac{d^2 y}{dx^2} + k^2 y = 0, y(0) = 0, y'(1) = 0])$			15	15	15	
111	$\text{solve}([\frac{dx}{dt} = x - y, \frac{dy}{dt} = x + y], [x, y])$			•	•	•	
112	verify the above is a solution			•	•	○	
113	$\text{solve}([\frac{dx}{dt} = x(1 + \frac{\cos t}{2 + \sin t}), \frac{dy}{dt} = x - y])$					○	
114	as above, but one equation at a time	○	○	○	•	○	•
115	$L = (D - 1)(D + 2)$ (operator)	•		•	•	•	
116	$L(f) \Rightarrow D^2 f + Df - 2f$	•		•	•	○	
117	$L_y(g(y)) \Rightarrow \frac{d^2 g}{dy^2} + \frac{dg}{dy} - 2g$	○		•	•	○	
118	$L_z(A \sin z^2)$	○		•	•	○	
119	$T = \sum_{k=0}^2 \frac{(D^k f)(a)}{k!} (x - a)^k$ (operator)	○		•	•	•	○
120	$T(f) \Rightarrow f(a) + (Df)(a)(x - a) + \dots$	•		•	•	•	○
121	$T_{y,b}(g(y)) \Rightarrow g(b) + \frac{dg}{dy} _{y=b}(y - b) + \dots$	•		•	•	•	○
122	$T_{z,c}(\sin z)$	•		•	•	•	•
123	compute Legendre polys directly	•	•	•	○	•	•
124	compute Legendre polys recursively	•	•	•	• ¹⁶	•	•
125	evaluate the 4 th Legendre poly at 1	○	○	•	•	○	○
126	$p = \sum_{i=1}^5 a_i x^i$	•	•	•	•	•	•
127	Horner's rule applied to the above			•	•	17	
128	convert to FORTRAN syntax	○	○	•	•	○	○
129	true and false \Rightarrow false	•	•	•	•	•	•
130	x or (not x) \Rightarrow true		•		•	•	•
131	x or y or (x and y) $\Rightarrow x$ or y		•		•		•